

PRX-T[®] lady

PROTOCOL FOR USE



REVITALISATION AND LIGHTENING OF EXTERNAL DELICATE AREAS

Frequency of sessions:

1 session every 15 days for 5 or more sessions.

METHOD OF USE

- 1) Clean the area to be treated with WiQomed **P SOLUTION**.
- 2) Perforate the cap of **PRX-T LADY** vial with a needle (18g).
- 3) Insert the syringe on the needle and extract the desired quantity.
- 4) Protect the mucous membranes with WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.
- 5) Apply (with syringe without needle) a limited quantity of product on the skin in the area being treated.
- 6) Apply with the rubber tipped brush covered with a nitrile glove, repeating the application 2-3 times.
- 7) Proceed area by area until the entire desired surface has been treated.
- 8) Remove the residue with a pad dampened with water.
- 9) In case of hyperpigmentation, continue the treatment with the home product WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS** twice a day before WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.
- 10) Restore the hydrolipidic film with WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.



LIGHTENING OF AXILLA

Frequenza sedute:

1 session every 15 days for 5 or more sessions.

METHOD OF USE

- 1) Clean the area to be treated with WiQomed **P SOLUTION**.
- 2) Perforate the cap of **PRX-T LADY** vial with a needle (18g).
- 3) Insert the syringe on the needle and extract the desired quantity.
- 4) Apply with the rubber tipped brush covered with a nitrile glove, repeating the application 2-3 times.
- 5) If irritation or intense burning occurs, stop the application.
- 6) Remove the residue with a pad dampened with water.
- 7) Apply WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS** and WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.

Continue the treatment with the home product WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS** and WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.



LIGHTENING OF MAMMARY AREOLA

Frequency of sessions:

1 session every 15 days for 5 or more sessions.

METHOD OF USE

- 1) Clean the area to be treated with WiQomed **P SOLUTION**.
- 2) Perforate the cap of **PRX-T LADY** vial with a needle (18g).
- 3) Insert the syringe on the needle and extract the desired quantity.
- 4) Apply on the areola with the rubber tipped brush covered with a nitrile glove, repeating the application 2-3 times (avoiding the nipple).
- 5) If irritation or an intense burning sensation occurs, stop the application.
- 6) Remove the residue with a pad dampened with water.
- 7) Apply WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS**.
- 8) Restore the hydrolipidic film with WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.



LIGHTENING OF PERIANAL AREA

Frequency of sessions:

1 session every 7 days for 5 or more sessions.

METHOD OF USE

- 1) Clean the area to be treated with WiQomed **P SOLUTION**.
- 2) Perforate the cap of **PRX-T LADY** vial with a needle (18g).
- 3) Insert the syringe on the needle and extract the desired quantity.
- 4) Protect the mucous membranes with WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.
- 5) Apply with the rubber tipped brush covered with a nitrile glove, repeating the application 2-3 times.
- 6) If irritation or an intense burning sensation occurs, stop the application.
- 7) Remove the residue with a pad dampened with water.
- 8) Apply WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS** and WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.

Continue the treatment with the home product WiQo **LIGHTENING SERUM FOR EXTERNAL DELICATE AREAS** and WiQo **RESTORING CREAM FOR EXTERNAL DELICATE AREAS**.



GENERAL CONTRAINDICATIONS IN THE APPLICATION OF PRX-T Lady

- Irritated skin
- Irritable skin
- Macerated skin
- Active dermatitis
- Allergy to kojic acid
- Allergy to latex, if latex gloves are used
- Pregnancy and breastfeeding
- Cutaneous and/or herpetic eruptions

WARNINGS

- Begin the application in a limited area. A slight tingling is normal. If the patient experiences intense burning, it means that the skin is irritable. In this case, rinse the treated area with water, stop the session and apply WiGo Restoring Cream for External Delicate Areas.
- Always apply the product on well lubricated skin **to avoid excoriations.**
- Carry out the treatment after a suitable time after shaving or waxing.
- In order to avoid allergic reactions, use nitrile rather than latex gloves.
- Do not treat areas affected by seborrhoeic dermatitis.
- **Do not treat the inguinal fold.**
- Warning: in case of contact with the eyes, rinse thoroughly with saline solution.

Possible side effects and problems:

- **Itchiness, with or without redness, in the days following the treatment:** very rare, presumably due to an allergy to kojic acid. Prescribe a steroidal cream.



Direct line to contact dermatologists and research biologists
for scientific questions or enquiries regarding outpatient treatment:
scientificdiv.gpq@wiqo.eu

GPQ srl

Operational Headquarters: Via Martinelli, 16 – 34015 Muggia (Trieste) - Italy
Tel.: +39 040 360408 - Fax: +39 040 761393 - gpq@wiqo.eu
www.wiqo.com